

amount to use is left to the expertise of the Veterinary Surgeon who may direct accordingly.  
As a guide: Cattle – up to 400ml.

**Contra-Indications, Warnings, etc**

Administer with care.  
The solution should be warmed to body temperature before use.  
Intravenous injections should be carried out slowly and stopped at once if untoward symptoms occur.  
When administered subcutaneously massage the injection site.

**Withdrawal Period**

Not indicated for milk or slaughter.

**Pharmaceutical Precautions**

Protect from light at a temperature not exceeding 25°C. This product is intended for single use only. Any product remaining after use should be discarded.  
Should any apparent growth/discolouration occur the product should be discarded.  
For animal use only. Keep out of reach of children.

**Legal Category** LM

**Package Quantities**

400ml amber glass vial.

VPA 10956/1/1/1

it is suggested that a dose of 100ml/500kg is given slowly intravenously followed by 100ml/500kg subcutaneously.

**Contra-Indications, Warnings, etc**

Administer with care particularly where concurrent toxæmia exists. The solution should be warmed to body temperature before use. Intravenous injections should be carried out slowly and stopped at once if untoward symptoms occur. When administered subcutaneously massage the injection site. Avoid the introduction of contamination during use.

**Withdrawal Period**

Meat: Nil; Milk: Nil.

**Pharmaceutical Precautions**

Protect from light at a temperature not exceeding 25°C. This product does not contain a preservative. Discard any unused solution. Should any apparent growth/discolouration occur the product should be discarded. For animal treatment only. Keep out of reach of children.

**Legal Category** LM

**Package Quantities**

100ml clear glass vial.

VPA 10956/1/1/1

**B.V.P. MEGACAL-M INJECTION  
(B.V.P. CALCIUM 50 PLUS INJECTION)**

**Presentation**

A clear, colourless to pale yellow, sterile injectable solution, containing calcium borogluconate and magnesium hypophosphite equivalent to 4.6% w/v calcium, 2% w/v phosphorus, 0.8% w/v magnesium.

**Uses**

The product is recommended for the treatment of cattle in the following conditions when complicated by magnesium and phosphorus deficiency.

- 1 Paresis caused by hypocalcaemia before, during and after birth, or during lactation.
- 2 Downer cow syndrome (including during pregnancy) caused by deficiency diseases or internal secretory dysfunction.

**Dosage & Route of Administration**

Administration by subcutaneous or slow intravenous injection using aseptic technique. The actual dose will depend upon the clinical condition of the animal and the amount to use is left to the expertise of the Veterinary Surgeon, who may direct accordingly. As a guide

**B.V.P. REPLENOLYTE (CONCENTRATE)**

**Presentation**

A clear, pink, aqueous, concentrated solution intended for dilution before use. For Oral use only.  
The product contains:

Calcium Chloride	0.5% w/v,
Potassium Chloride	1.5% w/v,
Magnesium Chloride	0.45% w/v,
Sodium Acetate	12.5% w/v,
Sodium Chloride	8.0% w/v,
Dextrose Monohydrate	50% w/v.

On reconstitution the available ion concentrations are:

Sodium	114mmol/L,
Potassium	10mmol/L,
Chloride	84mmol/L,
Dextrose	126mmol/L,
Magnesium	1.1mmol/L,
Calcium	1.7mmol/L,
Acetate	46mmol/L.

**Uses**

For Oral use to reverse the process of dehydration and electrolyte loss associated with scours in calves whether due to nutritional\* or infectious causes. To reduce scouring in bought-in calves. To reverse hypo-

glycaemia and aid recovery in cases of pregnancy toxæmia in ewes.

**Dosage & Route of Administration**

For oral administration only.

**Calves:**

Prior to use dilute one part of concentrate to twenty parts with water.  
As a guide:  
50ml concentrate to 1 litre with water.  
100ml concentrate to 2 litres with water.  
200ml concentrate to 4 litres with water.

**Scouring Calves:**

Withdraw all milk and milk replacer.  
Administer 2 litres of dilute Replenolyte twice daily for 2 days (four feeds). For the next 4 feeds use 1 litre of diluted Replenolyte and 1 litre of milk or milk replacer. Thereafter feed as normal.

**Sheep:**

In cases of pregnancy toxæmia in ewes 100ml of undiluted product should be administered in a suitable

drenching bottle. Treatment should be repeated hourly intervals as required.

**Contra-Indications, Warnings, etc**

In severe cases of dehydration additional intravascular rehydration therapy will be necessary. In such cases consult a veterinary surgeon. Ensure that ad colostrum is fed to animals.  
Do not hold diluted solution for longer than 24 h

**Withdrawal Period**

Nil.

**Pharmaceutical Precautions**

Store below 25°C. FOR ANIMAL TREATMENT C  
KEEP OUT OF REACH OF CHILDREN.

**Legal Category** LM

**Package Quantities**

1L, 2.5L, 5L plastic containers with aluminium foil and screw cap.

VPA 10956/1/7/1

f prior  
pend  
e con-  
if ther-  
n and  
serum  
dilation  
cularly  
er tox-  
e from  
andom  
see if  
copper  
ns can  
vell as  
s. It is  
nd sys-  
ernary  
1 cattle  
in one  
an be  
Avoid  
alfhood